

Description of the male of *Ctenophilothis altus* (Lewis, 1885): supplement to the revision of the genus *Ctenophilothis* Kryzhanovskij, 1987 (Coleoptera, Histeridae, Sapriniinae)

Tomáš Lackner¹

¹ Czech University of Life Sciences, Faculty of Forestry and Wood Sciences, Department of Forest Protection and Entomology, Kamýcká 1176, CZ-165 21 Praha 6 – Suchbát, Czech Republic

<http://zoobank.org/DB9F0BB4-BB76-496F-B7A0-A7991AE1AA6B>

Corresponding author: Tomáš Lackner (tomaslackner@me.com)

Received 11 August 2014
Accepted 9 September 2014
Published 24 October 2014

Academic editor:
Alexey Solodovnikov

Abstract

A description of the male, including illustrations of the male genitalia, and a colour image of the habitus of the extremely rare species *Ctenophilothis altus* (Lewis, 1885) are provided for the first time. Further support for the monophyly of the genus *Ctenophilothis* Kryzhanovskij, 1987 is given.

Key Words

Coleoptera
Histeridae
Sapriniinae
Ctenophilothis

In 2013 I published a revision of the genus *Ctenophilothis* Kryzhanovskij, 1987 (Lackner 2013). At the time of that revision this genus consisted of only two psammophilous Saharan species, both very rare. Although *C. chobauti* (Théry, 1900) has been found several times in the Moroccan and Algerian Sahara (Olexa 1990, Gomy et al. 2014), the other species, *C. altus* (Lewis, 1885) was known only from the female holotype and another female, collected in upper Egypt (Lackner 2013). During my recent visit to the Zoological Museum of the Humboldt University Berlin, Germany (ZMHUB) I discovered in the collection a male of this species, collected in Asyut, central Egypt.

In this short correspondence I publish the discovery of the third specimen of the extremely rare *C. altus*, together with the color image and illustration of its hitherto unknown male genitalia. Both Egyptian localities of this species are likewise mapped herein. As already noted by Lackner (2013) the genus *Ctenophilothis* is most probably monophyletic, sharing e.g. the absence of the antennal cavity, shortened or strongly reduced

lateral costa of the antennal groove or the peculiar shape of protibia (Lackner 2013: 281). The comparison of the male genitalia of both species reveals their striking similarities as well (compare figs 2–10 with those of Lackner 2013, figs 11–17) suggesting further support for the genus' monophyly. The main difference between the male terminalia lies in the shape of spiculum gastrale that in *C. chobauti* belongs to the most common type found in the Sapriniinae, having both 'head' and 'stem' sensu Caterino and Tishechkin (2013). On the other hand, the spiculum gastrale of *C. altus* is of a rather peculiar type, lacking typical 'head' or 'stem' (compare figs 7–8 with those of Lackner 2013, figs 14–15). Especially seen from the lateral view, the spiculum gastrale of *C. altus* is extremely flat, almost without projections (Fig. 8). According to my studies on the morphology of the Sapriniinae, the shape of the spiculum gastrale is very diverse and it was very difficult to parse this extraordinary diversity into discrete character states (Lackner in press).





Figure 1. *Ctenophilothis altus* (Lewis, 1885) habitus, dorsal view.



Figures 2–10. *Ctenophilothis altus* (Lewis, 1885) 2 – eighth sternite and tergite, ventral view; 3 – ditto, dorsal view; 4 – ditto, lateral view; 5 – ninth and tenth tergites, dorsal view; 6 – ditto, lateral view; 7 – spiculum gastrale, ventral view; 8 – ditto, lateral view; 9 – aedeagus, dorsal view; 10 – ditto, lateral view.




Figure 11. Distribution of *Ctenophilothis altus* (Lewis, 1885) in Egypt.

The second known locality of *C. altus* (Asyut, 27°11'00"N 31°10'00"E) lies more south than the single one published previously (Fig. 11). The presumed rarity of the species can be most likely ascribed by the insufficient collection aimed at the psammophilous Histeridae in Egypt. While efforts aimed at collecting psammophilous Sapriniinae of the Algerian and Moroccan Sahara yielded remarkable results (see e.g. Olexa 1990 or Gomy et al. 2014), there has been practically no such activity performed in Egypt.

Thanks are due to the Coleoptera curator of ZMHUB, B. Jaeger for his help and assistance during my visit to the collection as well as to Mahmoud Saleh Saleem (Riyadh, Saudi Arabia) for his help with the Egyptian localities. I would also like to thank the ZMHUB for the help with the open access of my work. This research received support from the SYNTHESYS Project <http://www.synthesys.info/>, which is financed by the European Community Research Infrastructure Action under the FP7 Integrating Activities Program as well as by the Internal Grant Agency (IGA n.20124364) Faculty of Forestry and Wood Sciences, Czech University of Life Sciences Prague, Czech Republic.

References

- Caterino MS, Tishechkin AK (2013) A systematic revision of *Operclipygus* Marseul (Coleoptera, Histeridae, Exosternini). *ZooKeys* 271: 1–401. doi: 10.3897/zookeys.271.4062
- Gomy Y, Labrique H, Francois A (2014) Contribution à la connaissance des Histeridae du Maroc (Coleoptera) (Troisième note) 1^{er} Partie. *Bulletin de la Société Entomologique de Mulhouse* 70(2): 23–35.
- Lackner T (2013) Revision of the genus *Ctenophilothis* Kryzhanovskij, 1987 (Coleoptera: Histeridae: Sapriniinae). *Zootaxa* 3691(2): 273–282. doi: 10.11646/zootaxa.3691.2.6
- Lackner T (in press) Phylogeny of the Sapriniinae subfamily reveals interesting ecological shifts in the history of the subfamily (Coleoptera: Histeridae). *Zoological Journal of the Linnean Society*.
- Olexa A (1990) On the genus *Philothis* and related genera (Coleoptera: Histeridae). *Acta Entomologica Bohemoslovaca* 87: 141–155.