Appendix 10 Pairwise genetic mean p distances (%) between the examined individuals of the European species of the Colletes succinctus group, based on the individuals investigated.

	P3	BK1	BK2	BK3	BK4	BK5	7346	7345	BK6	BK7	BK8	BK9	BK25	<b>BK26</b>	BK28	BK23	BK29	BK30	BK37	BK38
P3 (C. hederae)																				
BK1 (C. hederae)	0.32																			
BK2 (C. hederae)	0.64	0.95																		
BK3 (C. succinctus)	0.64	0.95	0.00																	
BK4 (C. succinctus)	0.64	0.95	0.00	0.00																
BK5 (C. succinctus)	0.64	0.95	0.00	0.00	0.00															
7346 (C. succinctus)	0.16	0.48	0.79	0.79	0.79	0.79														
7345 (C. succinctus)	0.16	0.48	0.79	0.79	0.79	0.79	0.32													
BK6 (C. collaris)	5.25	5.25	5.09	5.09	5.09	5.09	5.09	5.41												
BK7 (C. collaris)	5.09	5.09	4.93	4.93	4.93	4.93	4.93	5.25	0.16											
BK8 (C. collaris)	5.25	5.25	5.09	5.09	5.09	5.09	5.09	5.41	0.16	0.32										
BK9 (C. collaris)	5.25	5.25	5.09	5.09	5.09	5.09	5.09	5.41	0.16	0.32	0.00									
BK25 (C. pannonicus)	0.32	0.64	0.48	0.48	0.48	0.48	0.48	0.48	4.93	4.77	4.93	4.93								
BK26 (C. pannonicus)	0.48	0.79	0.48	0.48	0.48	0.48	0.32	0.64	4.77	4.61	4.77	4.77	0.48							
BK28 (C. brevigena)	0.00	0.32	0.64	0.64	0.64	0.64	0.16	0.16	5.25	5.09	5.25	5.25	0.32	0.48						
BK23 (C. brevigena)	0.32	0.64	0.64	0.64	0.64	0.64	0.48	0.48	5.25	5.09	5.25	5.25	0.64	0.48	0.32					
BK29 (C. brevigena)	0.64	0.95	0.00	0.00	0.00	0.00	0.79	0.79	5.09	4.93	5.09	5.09	0.48	0.48	0.64	0.64				
BK30 (C. brevigena)	0.00	0.32	0.64	0.64	0.64	0.64	0.16	0.16	5.25	5.09	5.25	5.25	0.32	0.48	0.00	0.32	0.64			
BK37 (C. brevigena)	0.32	0.32	0.95	0.95	0.95	0.95	0.48	0.48	5.25	5.09	5.25	5.25	0.64	0.79	0.32	0.64	0.95	0.32		
BK38 (C. brevigena)	0.16	0.48	0.48	0.48	0.48	0.48	0.32	0.32	5.09	4.93	5.09	5.09	0.16	0.32	0.16	0.48	0.48	0.16	0.48	
7348 (C. creticus)	6.84	6.84	6.84	6.84	6.84	6.84	7.00	7.00	7.95	7.79	7.95	7.95	6.84	6.68	6.84	6.84	6.84	6.84	6.68	6.68